

Die Fortsetzbarkeit von Differentialformen auf
arithmetischen Quotienten von hermiteschen symmetrischen
Räumen
– Abstract –

Klaus Pommerening

J. reine angew. Math. 356 (1985), 194-220

Let D be a hermitean symmetric space of noncompact type, hence isomorphic to a bounded symmetric domain. Let a discontinuous group Γ act on D , and let Γ be arithmetically defined. Let X_Γ be the BAILY-BOREL compactification of D/Γ and X , a desingularization of X_Γ . Let $A[\Gamma, \wedge^p]$ be the space of Γ -automorphic forms for the natural automorphy factor that belongs to the p -forms and $\Omega^p(D)^\Gamma$, the space of Γ -invariant p -forms on D .

Then we have inclusions

$$\Omega^p(X) \hookrightarrow A[\Gamma, \wedge^p] \hookrightarrow \Omega^p(D)^\Gamma.$$

The equality at the second arrow is the question of KOECHER's principle. The equality at the first arrow is the main result of this paper: Every Γ -automorphic p -form on D of degree $p < \dim D$ extends to X .

The analogous result for $p = \dim D$ is false; here only the cusp forms extend [1]. The special case where D is SIEGEL's half-space and Γ is commensurable with SIEGEL's modular group was treated in [2].

References

- [1] Y.-S. Tai in: A. Ash, D. Mumford, M. Rapaport, Y.-S. Tai, *Smooth Compactification of Locally Symmetric Varieties* Brookline 1975.
- [2] E. Freitag, K. Pommerening; J. reine angew. Math 331 (1982), 207–220.